

## Observation of Barrett's esophagus and early gastric cancer by magnifying Narrow Band Imaging (NBI) endoscopy

T. Endo

Sapporo Shirakabada Hospital, Sapporo, Japan

By only changing the optical filters for sequential lighting from the conventional broadband type to the narrow band type, NBI system can be installed. Through our clinical evaluations as to Barrett's mucosa and early gastric cancers, we confirmed NBI has an advantage over a conventional system with the representation of the important observation and diagnosis such as the fine mucosal patterns and capillary structure. We believe NBI will be one of the tools supporting the endoscopic diagnosis of Barrett esophagus and tumor related neo angiogenesis of early gastric cancers.

In this workshop, Firstly I would like to report the diagnostic implication of fine mucosal patterns (pit diagnosis) obtained by NBI magnifying pictures *in vivo* compared with conventional and chromo-endoscopy with reference to the discrimination of SCE from the remaining epithelial types. The relationship between the endoscopic and histopathologic diagnosis was more accurate by NBI endoscopy than conventional magnifying endoscopy.

Secondly I would like to show the precise observation of the esophago-gastric junction as to the discrimination among epithelial types such as the squamous, cardiac and fundic mucosa using NBI system.

Finally depressed type early gastric cancers were observed by NBI. The fine mucosal vascular pattern was recorded and compared with the histological differentiation and features of vessels by staining anti-CD34 antibody. In some patients, c DNA array analysis was performed as to differences among histological types. Tumor vascular patterns were classified into two categories. Grid like network patterns not only characterized differentiated type but were also associated with high micro vascular density. Short twig-like patterns typified the undifferentiated type and a low vascular density. Differentiated types highly expressed some angiogenic factors, such as VEGF and Flt-4.

## Can we make a diagnosis of *Helicobacter pylori* infection endoscopically?

R. Narisawa<sup>1</sup>, Y. Satoh<sup>2</sup>, M. Kobayashi<sup>2</sup>, M. Takeuchi<sup>1</sup>,  
J. Yokoyama<sup>1</sup>, K. Shioji<sup>1</sup>, K. Sugimura<sup>2</sup>, U. Aoyagi<sup>2</sup>

<sup>1</sup>Department of Endoscopy, Niigata University Medical and Dental Hospital

<sup>2</sup>Division of Gastroenterology and Hepatology, Niigata University, Graduate School of Medical and Dental Sciences, Niigata, Japan

Since *Helicobacter pylori* (*H. pylori*) had been discovered in 1983, the diagnosis and treatment of upper gastrointestinal diseases have changed greatly.

Today, there have been many reports that *H. pylori* related to many diseases of gastrointestinal tract. As a result, the diseases that considerable cause was regarded as *H. pylori* infection were as follows: gastric ulcer, duodenal ulcer, MALT lymphoma of the stomach, acute gastric mucosal lesion (AGML), chronic gastritis (especially, chronic active gastritis, atrophic gastritis,

superficial gastritis, and follicular gastritis), gastric cancer, gastric adenoma, hyperplastic polyp of the stomach, giant rugae of the stomach, and so on. However, the explication of the detailed mechanism between contracting such diseases and *H. pylori* infection was not so clear even now.

On the other hand, most of fundic gland polyps of the stomach were found out in patients with *H. pylori* negative, and most of Kammroetung (Comb-like redness) of the stomach was also. And RAC (regular arrangement